

Prevalence of Depression Among Community Dwelling Elderly in Karachi, Pakistan

Sayed Muhammed Mubeen, MBBS, MES, MCPS, MPH* , Danish Henry, MBBS*
Sarah Nazimuddin Qureshi, MBBS*

(Received: 1 Dec 2012; Revised: 14 Jun 2012; Accepted: 24 Jul 2012)

Objective: The objectives of the study were to find out the prevalence of depression and to identify associated risk factors among community dwelling elderly in Karachi.

Methods: It was a cross-sectional, descriptive study involving 284 community-dwelling elderly residing in Karachi, Pakistan. A non-probability convenience sampling was done. The Geriatric Depression Scale (GDS-15) was used to assess depression. Descriptive statistics was performed using SPSS version 12. Cross tabulation for different variables was done and Chi-square was used as test of significance. The level of significance was set as $p < 0.05$. An informal (verbal) consent was taken. Anonymity and confidentiality was assured.

Results: Among 284 respondents, 74% were males while 26% were females. The mean age was 68.44 ± 7.59 years. The study found that 16.5% respondents were depressed while 23.6% were suggestive of depression. Depression was more among men than in women. Depression was statistically significant among married respondents ($p < 0.05$) and illiterate ($p < 0.001$). Although a large proportion of the participants were satisfied with their income, this was statistically significant ($p < 0.001$) for depression among those who were not satisfied with their income. Similarly, sleep was significantly disturbed ($p < 0.001$) among the depressed respondents.

Conclusion: A significant prevalence of geriatric depression was reported. In order to reduce its prevalence, general physicians and other health care professionals need to be sensitized about geriatric depression and its risk factors.

Declaration of interest: None

Citation: Mubeen SM, Henry D, Nazimuddin Qureshi S. Prevalence of depression among community dwelling elderly in Karachi, Pakistan. *Iran J Psychiatry Behav Sci* 2012; 6(2): 84-90.

Keywords: ●Aged ●Depression ●Geriatrics

Introduction

Depression is a common psychiatric disorder and the most common in geriatrics (1). Currently, depression is the 2nd leading cause of Disability Adjusted Life Years (DALYs) in the age category 15-44 years for both sexes combined and by the year 2020, it is projected to reach the 2nd place of the

ranking of DALYs, calculated for all ages, both sexes (2).

Various studies have been conducted to investigate depression in the elderly and many found depression to be largely under diagnosed and untreated (3). Primary-care doctors rarely diagnose depression and, when they do provide inappropriate treatment (4). Barriers to adequate diagnosis and treatment include doctors' reluctance to discuss emotional problems, time constraints and medical co-morbidities, complicating diagnosis and competing for medical attention. Perceived stigma contributes to patients' reluctance to initiate psychiatric treatment (5). While late-life depression tends

Authors' affiliation: * Associate Professor, Hamdard College of Medicine Dentistry, Hamdard University, Karachi, Pakistan.

●**Corresponding author:** Dr. Syed Muhammed Mubeen, Hamdard College of Medicine Dentistry, Madinat al-Hikmah, Mohammed Bin Qasim Ave, Hamdard University, Karachi, 74600, Pakistan.
Tel : 009221-36440091
Fax : 009221-36440091
E-mail: dr_mubeen@hotmail.com

to be a recurring disorder, it may be masked by hypochondriasis or somatization (6). Furthermore, complex medication regimens undermine treatment adherence. It is not surprising that older patients with depressive disorder have high rates of treatment resistance and mortality (6,7).

In this globalized world, the 'Empty Nest Syndrome' is not just a Western phenomena; the younger generation has allowed it to make in-roads into the eastern culture. In Pakistani social setting, many people work past their retirement age (60 years in Pakistan) mainly due to compelling circumstances to make both ends meet. Also, the gradual disintegration of the centuries old joint family system, emergence of nuclear family system and individuality – self centeredness has eroded the channels of care for the elderly and has banished them to isolation with impunity (8).

Contributing factors include a fragile health system with no dedicated geriatric clinics, deteriorating social and moral fiber add to miseries of the elderly with poor checks and balances in pensions and old-age benefits, mechanical and fast way of life in the urban areas. It is coupled with intolerance, lack of harmony with nature, rat race for money and power, ruthlessness, disposable culture and rampant consumerism, looming insecurity, social injustice all are likely to drag a person into the gallows of depression, sometimes never to come back. Being denied what they have earned throughout their lives, these senior citizens become more prone to depression.

The unattended depression is further aggravated by health care professional's inability to recognize the importance of mental health. A study done in Lahore, Pakistan revealed that 50% of the medical students and professionals claimed that they had not heard about depression and a significant proportion of all the people surveyed had a negative attitude towards depressed patients (5). The objectives of the study were to find out the prevalence of

depression among community-dwelling elderly in Karachi.

Materials and Methods

It was a cross-sectional, descriptive study using non-probability convenience sampling method spanned over a period of 12 months (March 2009 to February 2010). The sample included community-dwelling elderly population residing in one of the four districts of Karachi, Pakistan. As there is no general agreement on the age at which a person becomes old, most of the developed world countries have accepted the chronological age of 65 years as a definition of 'elderly' or older person. Moreover, there is no UN standard numerical criterion, but the UN agreed cut-off is >60 years to refer to the older population (9). Considering the average lifetime expectancy of 67 years among a Pakistani population (10), elderly in this study refers to persons of age 60 years and above. With an estimated prevalence of 25% reported in pervious studies with 95% level of confidence and an absolute precision of 0.05, a sample size of 289 was calculated. Nonetheless, 300 persons were included in the study. Questionnaire of sixteen respondents were not included due to incomplete information, hence, 284 respondents were included as the final sample size. Elderly persons regularly visiting parks in afternoons and evenings and attending Masjid and church (places of worship) were invited to participate in the study. People going independently to these places are likely to have attentive cognition and can thus are able to sensibly respond to the survey questionnaire.

The questionnaire used in the study was divided into socio-demographic information and the GDS-15. The socio-demographic information included sex, age, residence, religion, mother tongue, education, marital status, occupation and family setup. It comprises Joint Family System (JFS) having two or more nuclear families that form a corporate economic unit (11) and Nuclear

Family System (NFS) consisting of parent and their dependent children (11). Co-morbidities, use of daily medications and addiction(s) were also inquired.

The second part comprised of a scale for assessing late-life depression. It refers to depressive syndromes defined in the DSM-IV and in the ICD-10 that arise in adults older than age 65 years (3). For this assessment, The GDS-15, the shorter version of the questionnaire was selected. The usage of GDS-15 can be attributed to its easy administration to the target population and concise content, taking 5 to 7 minutes to complete. The scale can be used in healthy, medically ill and mild to moderate cognitively impaired older adults (12). This scale has been used both as a self-administered questionnaire and as read out interviews.

The GDS-15 assesses depression in the elderly on a score of 15. Scores of 0-4 are considered normal, depending on age, education, and complaints; 5-8 indicate mild depression; 9-11 signifies moderate depression and 12-15 specifies severe depression. Any positive score above 5 on the GDS-15 should prompt an in-depth psychological assessment and evaluation for suicidality (12).

The GDS was found to have 92% sensitivity and 89% specificity when evaluated against diagnostic criteria. The validity and reliability of the tool have been supported through both clinical practice and research (13,14). Validation studies of the GDS-15 established it as a promising screening device for detecting depression in the elderly. An Australian study found GDS-15 to be a good screening instrument for major depression as defined both by ICD-10 and DSM-IV (15). Similarly, GDS with 10 and 15 items produced consistent results in the assessment of elderly patients when total scores were used as clinical guidelines (15). Nevertheless, GDS does not assess suicidality and is not a substitute for a diagnostic interview by a mental health professional (12).

The inclusion criteria were consenting individuals above the age of 60 years

irrespective of sex, ethnicity or religion. Elderly suffering from chronic diseases like diabetes and hypertension were also included. Individuals below the age of 60, elderly suffering from known dementia and institutionalised elderly (in hospitals, old age homes, etc.) were excluded from the study.

Statistical Package for Social Sciences (SPSS - V12) was used for statistical analysis and data management. The data was serialized for facilitating computer entries. Descriptive statistics was performed. Cross tabulation for different variables was done and Chi-square was used as test of significance to find out the association between two variables. The level of significance was set as p less than 0.05.

The respondents of the questionnaire were informed of the purpose of the study and the usage of the information they provided. An informal (verbal) consent was taken before the respective respondents filled the questionnaire. The researcher(s) read out the questions for the respondents unable to read and write and recorded the responses. Anonymity and confidentiality of their information was assured and the right to withdraw upheld. The Research Committee of Hamdard University, Karachi gave the ethical approval.

Results

Among 284 respondents, males (74%) were in a numerical majority than females (26%) and nearly half (47%) of the respondents were in the age bracket of 60 to 65 years. The mean age was 68.44 ± 7.59 years. The majorities were Muslims (90%), married (77%) and spoke Urdu (72.4%) language. A little more than a quarter was illiterate and 7% of the respondents were unemployed at the time of study. Sixty-two percent were living as joint families while 47% of the respondents lived with their children only. Self-income was generated by 114 (40%) of the respondents whereas 55 (19%) rely on pension only. However, the majority (71%) were found to be satisfied with their income (Table 1).

Table 1. Demographic profile of the community dwelling elderly in Karachi Pakistan

	N	%
Sex		
Male	210	73.9
Female	74	26.1
Age (in years)		
60 – 65	132	46.5
66 – 70	69	24.3
71 – 75	39	13.7
76 – 80	28	9.9
> 80	16	5.6
Marital status		
Single	12	4.2
Married	218	76.8
Widow	43	15.1
Separated / Divorced	11	3.9
Religion		
Islam	256	90.1
Hinduism	19	6.7
Christianity	9	3.2
Educational status		
Primary	47	16.5
Secondary	67	23.6
Intermediate	39	13.7
Graduate/diploma	32	11.3
Postgraduate	19	6.7
Illiterate	80	28.2
Occupational status		
Employed / self-employed	96	33.8
Retired	57	20.1
Home maker/Unemployed	20	7.0
Mother tongue		
Urdu	207	72.4
Punjabi	29	10.2
Sindhi	22	7.7
Pushto	21	7.4
Baluchi	5	1.8
Family system		
Joint	175	61.6
Nuclear	109	38.4
Living status		
Alone	34	12.0
With spouse only	57	20.0
With children only	134	47.2
Both spouse and children	59	20.8
Source of income		
Self	114	40.1
Children	76	26.8
Pension	55	19.4
Savings	34	12.0
Charity	5	1.8
Income satisfaction		
Yes	203	71.5
No	81	28.5

Figure 1 highlighted the presence of depression among the 284 respondents. It was found that 47 (16.5%) respondents were depressed while 67 (23.6%) were suggestive of depression. Depression was mainly present in persons in the age range of 60-70 years.

When demographic variables were cross-tabulated with mental status, depression was more among men than in women ($p < 0.05$).

Similarly, depression was found to be statistically significant among married respondents ($p < 0.05$) and those who were illiterate ($p < 0.001$). Although a large proportion of the participants were satisfied with their income, this was found to be statistically significant ($p < 0.001$) for depression among those who were not satisfied. Similarly, sleep was significantly disturbed ($p < 0.001$) among the depressed participants. (See table 2).




Figure 1. Geriatric Depression Scale (GDS) Scores of the community dwelling elderly in Karachi, Pakistan

Discussion

An individual strives throughout his or her life for a peaceful old age, financial security, health, kinship and independence and yet these things remain elusive to many on this planet. The feelings of sadness that accompany routine life are usually appropriate and transitory, and can even present an opportunity for personal growth. However, when depression persists and impairs daily life, it may be an indication of a depressive disorder.

The present study reported 16.5% of the elderly population to be depressed whereas two hospital-based studies in Karachi showed depression in 23% and 20% among the elderly population (11,16). A systemic review revealed that a prevalence of 34% for anxiety and depressive disorders in Pakistani population (17). Moreover, there is a wide range (10% to 60%) of prevalence of depression in other locally published studies in Pakistan. These figures can be compared with neighboring

country India having similar socio-demographic structure, the prevalence rates for depression showed a variation between 13% to 31% (18,19). A recent study done in Iran reported 23.5% of the subjects suffered from depression (20).

The relatively low prevalence of depression in this study can be attributed to the fact that the study was conducted in community dwelling elderly while previous studies conducted among elderly population of Karachi were based in a tertiary care setting. Moreover, elderly individuals coming to a tertiary care setting

Table 2. Cross tabulation of demographic variables and mental status of community dwelling elderly in Karachi, Pakistan

Variables	Mental Status		p-value†
	Normal (%)	Suggestive of depression / Depressed (%)	
Sex			
Male	116 (55.2)	94 (44.8)	0.04
Female	51 (68.9)	23 (31.1)	
Age (in years)			
60 – 70	113 (56.2)	88 (43.8)	NS
71 – 80	44 (65.7)	23 (34.3)	
> 80	10 (62.5)	6 (37.5)	
Marital status			
Single	2 (16.7)	10 (83.3)	0.02
Married	134 (61.5)	84 (38.5)	
Widow	27 (62.8)	16 (37.2)	
Separated / Divorced	7 (63.6)	4 (36.4)	
Religion			
Islam	150 (58.6)	63 (24.6)	NS
Others	20 (71.4)	4 (14.3)	
Educational status			
up to Intermediate	81 (52.9)	41 (26.8)	< 0.001
up to Graduate/diploma	43 (84.3)	6 (11.8)	
Illiterate	46 (57.5)	20 (25)	
Occupational status			
Working	66 (58.6)	45 (41.4)	NS
Retired/Housewife/ Unemployed	104 (59)	69 (41.0)	
Mother tongue			
Mohajir (Urdu speaking)	129 (62.3)	78 (37.7)	NS
Others	41 (53.2)	36 (46.8)	
Family system			
Joint	106 (60.6)	69 (39.4)	NS
Nuclear	64 (58.7)	45 (41.3)	
Living status			
Alone	15 (44.1)	19 (55.9)	0.03
With spouse only	20 (50.9)	28 (49.1)	
With children only	90 (62.2)	44 (32.8)	
Both spouse and children	36 (61.0)	23 (39.0)	
Source of income			
Self	64 (54.1)	50 (43.9)	NS
Any other means	106 (62.4)	64 (37.6)	
Income satisfaction			
Yes	152 (74.9)	51 (25.1)	< 0.001
No	15 (18.5)	65 (81.5)	
Sleep disturbances			
Yes	60 (47.2)	67 (52.8)	< 0.001
No	107 (68.2)	50 (31.8)	

†Variables were combined for statistical significance.

certainly present with health problems (the concept of wellness clinics is not widely prevalent in Pakistan) making them more prone to depression. In the community-dwelling individuals, normal elderly with controlled co-morbidities or no co-morbidities were also included from all socio-economic classes. Moreover, depression was mainly found in the 60-65 year age bracket indicating that a sizeable elderly population is in this age group. This is supported by UNICEF statistics (2009) which shows the average life expectancy of a Pakistani to 65 years (10).

In our study, depression was found to be mainly affecting males. A Brazilian study concluded that being male was a risk factor for depressive symptoms (21). On the other hand, studies have shown that women suffer from depression more than men do (16, 20, 22-24). There is a possibility that women seek treatment for depression while men are reluctant to accept their depressive state as they are the bread earners and are more prone to face the harsh realities of the world. However, in the present study, there were more male respondents than females (around ¾ of the sample). Further studies involving equal males and females can confirm this sex predilection.

In our study, those who were depressed, more individuals were residing in a joint family than in a nuclear family. Contrary to this, a Karachi based study showed that elderly living in a nuclear family system were 4.3 times more likely to suffer from depression than those living in a joint family system (11). This shift indicates that individuals residing in a joint family are exposed to new challenges, possibly of augmented generation gap and their inability to cope up with the ever-changing world. Conversely, individuals in a nuclear family do things their way and therefore escape the clashes that emerge when an experienced mind interact with the youthful, energetic, risk-taking young individual seeking pace with an elusive materialistic world.

In the present study, 72% of the respondents were satisfied with their income. However, among those who were not satisfied, 82% were found to be depressed. These results were supported by other studies that showed

significant differences between depression and poor financial status in elderly population (20,25). Hence, income satisfaction is central to depression in the elderly. Elderly, mostly retired find it extremely difficult to survive in financial uncertainty and rising costs of health care. A similar result was found in a study conducted in the US where found the incidence and persistence of depression to be higher among persons with low incomes (26). The unfulfilled economic needs bring frustration and certainly disappointment.

Limitation

The presence of depression warrants prompt intervention and treatment. Any positive scores above five on the GDS in the present study prompt an in-depth psychological assessment and evaluation for suicidality. However, this remains to be one of the limitations of the study due to the lack of follow-up by the researchers.

Conclusion

The prevalence of geriatric depression reported by our study is less than that most of the other countries but still it is a significant percentage. If concerted and continuous efforts are endeavored by both health care professionals and the government, a reduction in the prevalence of geriatric depression is possible. As there are no specialized geriatric clinics in Pakistan, general physicians and other health care professionals need to be sensitized about geriatric depression, its risk factors and should treat it. Further, population-based studies should be carried out to estimate the prevalence of geriatric depression throughout Pakistan, explore risk factors and their relationship to depression and figure out potential solutions.

Authors' contributions

SMM conceived and designed the study, performed the statistical analysis, and helped to draft the manuscript and its critical final revision. DH carried out data collection, contributed in statistical analysis and drafted the study design evaluation and results. SNQ

re-evaluated and interpreted the data, drafted and revised the manuscript. All authors read and approved the final manuscript before submission to the journal.

References

1. Adamek ME, Slater GY. Depression and Anxiety. *J Gerontol Soc Work* 2008;50(1):153-89.
2. WHO. Depression. Geneva: World Health Organization; 2009 [cited 2009 October 06]; Available from: URL: http://www.who.int/mental_health/management/depression/definition/en/.
3. Alexopoulos GS. Depression in the elderly. *Lancet*. 2005; 365:1962.
4. Bruce ML, McAvay GJ, Raue PJ, Brown EL, Meyers BS, Keohane DJ, et al. Major depression in elderly home health care patients. *Am J Psychiatry* 2002;159(8):1367-74.
5. Naeem F, Ayub M, Izhar N, Javed Z, Irfan M, Haral F, et al. Stigma and Knowledge of Depression: A survey comparing medical and non-medical students and staff in Lahore, Pakistan. *Pak J Med Sci* 2005;21(2):155-8.
6. Casey DA. Depression in the elderly. *South Med J*. 1994;87(5):559-63.
7. Spear J, Chawla S, O'Reilly M, Rock D. Does the HoNOS 65+ meet the criteria for a clinical outcome indicator for mental health services for older people? *Int J Geriatr Psychiatry* 2002;17(3):226-30.
8. Itrat A, Taqui AM, Qazi F, Qidwai W. Family systems: perceptions of elderly patients and their attendants presenting at a university hospital in Karachi, Pakistan. *J Pak Med Assoc* 2007;57(2):106-10.
9. WHO. Definition of an older or elderly person. World Health Organization; 2009 [cited 2009 October 30]; Available from: URL: www.who.int/entity/healthinfo/survey/ageingdefnolder/en/.
10. UNICEF. Pakistan: Demographic indicators. 2009 [updated October]; Available from: URL:

- http://www.unicef.org/infobycountry/pakistan_pakistan_statistics.html#78.
11. Taqui A, Itrat A, Qidwai W, Qadri Z. Depression in the elderly: Does family system play a role? A cross-sectional study. *BMC Psychiatry* 2007;7:57
 12. Kurlowicz L, Greenberg SA. The Geriatric Depression Scale (GDS). New York: The Hartford Institute of Geriatric Nursing, College of Nursing; 2007; Available from: URL: http://consultgerirn.org/uploads/File/trythis/try_this_4.pdf
 13. Mitchell AJ, Bird V, Rizzo M, Meader N. Diagnostic validity and added value of the Geriatric Depression Scale for depression in primary care: a meta-analysis of GDS30 and GDS15. *J Affect Disord* 2010;125(1-3):10-7.
 14. Wancata J, Alexandrowicz R, Marquart B, Weiss M, Friedrich F. The criterion validity of the Geriatric Depression Scale: a systematic review. *Acta Psychiatr Scand* 2006;114(6):398-410.
 15. Almeida OP, Almeida SA. Short versions of the geriatric depression scale: a study of their validity for the diagnosis of a major depressive episode according to ICD-10 and DSM-IV. *Int J Geriatr Psychiatry* 1999;14(10):858-65.
 16. Ganatra HA, Zafar SN, Qidwai W, Rozi S. Prevalence and predictors of depression among an elderly population of Pakistan. *Aging Ment Health* 2008;12(3):349-56.
 17. Mirza I, Jenkins R. Risk factors, prevalence, and treatment of anxiety and depressive disorders in Pakistan: systematic review. *Bmj* 2004;328(7443):794.
 18. Barua A, Kar N. Screening for depression in elderly Indian population. *Indian J Psychiatry* 2010; 52(2): 150-3.
 19. Kamble SV, Dhumale GB, goyal RC, Phalke DB, Ghodke YD. Depression among elderly persons in a primary health centre area in Ahmednagar, Maharashtra. *Indian J Public Health* 2009;53(4):253-5.
 20. Majdi MR, Mobarhan MG, Salek M, Taghi M, Mokhber N. Prevalence of Depression in an Elderly Population: A Population-Based Study in Iran. *Iran J Psychiatry Behav Sci* 2011;5(1):17-21.
 21. Marinho PE, Melo KP, Apolinario AD, Bezerra E, Freitas J, Melo DM, et al. Undertreatment of depressive symptomatology in the elderly living in long stay institutions (LSIs) and in the community in Brazil. *Arch Gerontol Geriatr* 2010;50(2):151-5.
 22. Ostbye T, Kristjansson B, Hill G, Newman SC, Brouwer RN, McDowell I. Prevalence and predictors of depression in elderly Canadians: The Canadian Study of Health and Aging. *Chronic Dis Can* 2005;26(4):93-9.
 23. Chan A, Malhotra C, Malhotra R, Ostbye T. Living arrangements, social networks and depressive symptoms among older men and women in Singapore. *Int J Geriatr Psychiatry* 2011;26(6):630-9.
 24. Malhotra R, Chan A, Ostbye T. Prevalence and correlates of clinically significant depressive symptoms among elderly people in Sri Lanka: findings from a national survey. *Int Psychogeriatr* 2010;22(2):227-36.
 25. Chou KL, Chi I, Chow NW. Sources of income and depression in elderly Hong Kong Chinese: mediating and moderating effects of social support and financial strain. *Aging Ment Health* 2004;8(3):212-21.
 26. Zimmerman FJ, Katon W. Socioeconomic status, depression disparities, and financial strain: what lies behind the income-depression relationship? *Health Econ*. 2005;14(12):1197-215.