

Comparison Between the Mental Health of Mothers of Children With Autism and Control Group

Forough Riahi, MD* , Sakineh Izadi-mazidi, MSc**

(Received: 21 Jun 2010, Revised: 14 May 2012, Accepted: 9 Jun 2012)

Objective: The aim of this study was to compare the mental health of mothers of children with autism and those of control group.

Methods: Sample of this study consists of 32 mothers of Children with Autistic disorder and 29 mothers of children without Autism; referring to hospitals of Ahvaz city using Convenience sampling. All subjects were asked to complete Demographic questionnaire and General Health Questionnaire (GHQ). Data were analyzed using Multivariate Analysis of variance (MANOVA).

Results: Results indicated that significant differences exist in mental health ($F=5.69$, $P=0.02$) and anxiety/insomnia ($F=4.82$, $P=0.03$), between mothers of children with Autism and control group. There were not any other significant differences in the other subscales.

Conclusion: It is essential using some mental health improving interventions for mothers of children with Autistic disorder.

Declaration of interest: None.

Citation: Riahi F, Izadi S. Comparison between the mental health of mothers of children with autism and control group. *Iran J Psychiatry Behav Sci* 2012; 6(2): 91-95.

Keywords: •Autism •Children •Mental health •Mother

Introduction

Parents' mental health could be intensively influenced by disabled child, especially when there is a disorder such as Autism including a wide range of behaviors and particularly social behaviors (1). Autistic disorder is one of the pervasive developmental disorders characterized by delay and deviance in the development of social, communicative and other skills, various motor mannerisms, resistance to change, and idiosyncratic interests and preoccupations (2). Existing studies suggest that characteristics of the disorder cause stress in parents. And the more severe the child's

symptoms, the greater will be the degree of parental stress (3). And some studies found that they often experience frustration and pessimism, and they have particular characteristics such as schizoid traits, higher traits of aloof, hypersensitive, anxious, tense, and rigid which in turn contribute to their increased psychological stress (4,5). It also has been reported that parents of children with autism experience more marital distress and conflicts with non-disabled children (6). Sharply et al state that mothers experience greater impact than fathers (4). They blame themselves for their children's disorder, more likely than mothers of normative children (6). They are also the parent who was most likely to be held responsible for their child's behavior (7).

Having an autistic child has a remarkable effect on mother's careers. Those who are able to work are often forced to miss work, performing below their normal level or getting back to part-time status. Resentment is

Authors' affiliations: *Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. **Shahid Chamran University, Ahvaz, Iran

•**Corresponding author:** Sakineh Izadi, Graduate student of clinical psychology, Ahvaz Shahid Chamran University, Ahvaz, Iran.
Tel: +989171346040
Fax: +986113743038
E-mail: S.izadi.m@gmail.com

observed among mothers caused by this problem (3).

So it is necessary to study mother's problems and their psychological needs (because mothers are the family members with the most connection with these children). More knowledge about the adaptation process and development of psychopathologies in parents will let us design more effective strategies and programs to help them (5). But few studies have been performed in Iran (1). The purpose of this study was to compare the mental health of mothers of children with Autism to mothers of children without autism.

Materials and Methods

Sample of this Study included 32 mothers with autistic children who referred to child and adolescent psychiatrist, and 29 mothers without children autism referred to pediatrician in Ahvaz city hospitals. Two groups were selected by convenience sampling and were matched based on marital status, occupational status, level of education and ethnicity (Table 1). Children with autistic disorder were diagnosed by a child and adolescent psychiatrist, according to DSMIV-TR criteria. And children without autism didn't suffer from any physical or mental illness.

Data was gathered using GHQ. The questionnaire presented by Goldberg and Hillary for screening of non-psychotic psychological disorders, in 1979. It has 28 questions and four sub-scales and each scale has 7 questions. Adham et al evaluated validity of test 91% and 88% respectively. Reliability was calculated by Cronbach's alpha as a measure of internal consistency reliability. Cronbach's alpha was 0.84 for physical symptoms, social functioning 79%, depression 81% and mental health 91% (8).

Noorbala, Bagheriyazdi, Mohammad (2009) evaluated the concurrent validity of Persian form of the questionnaire by the Symptom Checklist- 90- Revised. They found that there was a significant correlation between scores of samples within the

subscales of depression ($r=0.72$), Anxiety ($r=0.75$), physical symptoms ($r=0.67$) and total score ($r=0.84$) of these instruments ($p<0.001$). The evaluation of reliability of GHQ-28 carried out by the test-retest procedure a week after the first stage. The correlation was significant ($r=0.85$, $p<0.01$) (9). Demographic data questionnaire was also filled in.

Data were analyzed using MANOVA. The probability level of 0.05 was accepted statistically significant. Statistical analyses were carried out by SPSS software, version 16.

Results

Mean (\pm SD) age of mothers with and without autistic children were 34.12 (\pm 5.9) and 37.27 (\pm 5.2) respectively ($p=0.09$). Frequencies of the mothers' demographic features are listed in table 1.

Table 1. Frequencies of the mothers' demographic features

Feature	Frequency		P-value
	mothers with autistic children	mothers without autistic children	
Marital status			
Married	32(100%)	29(100%)	-
Divorced/widowed	0	0	
Occupation status			
Yes	2(7%)	1(3%)	0.62
No	30(93%)	28(97%)	
Level of education			
High school	6(19%)	7(24%)	0.33
Diploma	19(59%)	17(59%)	
Bachelor	6(19%)	4(14%)	
Post graduate	1(3%)	1(3%)	
Ethnicity			
Fars	16(50%)	18(62%)	0.43
Arab	7(22%)	5(17%)	
Lor	9(28%)	6(21%)	
Other	0	0	

Tables 2, shows the mean and Standard deviation of mental health variable and its subscales in mothers with and without autistic children; and results of MANOVA.

As shown in table 2, there are significant differences between mothers with and without autistic children in mental health variable ($F=5.69$, $P=0.02$) and its subscale, anxiety/

insomnia ($F=4.82$, $P=0.03$). There are not any other significant differences in the other subscales.

normal lifestyle, lifetime care, worrying about the future of children and blaming mother as being responsible for disorder, threat mental health of family members and cause anxiety, especially in mothers(3,14-16).

Table2. Mean and SD of mental health variable and its subscales in mothers with and without autistic children; and results of MANOVA

Variable	Mean(\pm SD)		df	Mean Square	F	P-value
	mothers with autistic children	mothers without autistic children				
Mental health	58.71 \pm 13.29	51.34 \pm 10.51	1	827.20	5.69	0.02
physical symptoms	13.96 \pm 4.40	13.20 \pm 3.74	1	8.83	0.52	0.47
Anxiety/insomnia	16.78 \pm 8.21	13.03 \pm 4.30	1	213.56	4.82	0.03
social dysfunction	17.62 \pm 2.56	16.78 \pm 3.05	1	10.72	1.33	0.25
Depression	15.65 \pm 5.31	14.68 \pm 5.05	1	14.21	0.52	0.47

Discussion

Results indicated that mothers of children diagnosed with Autism had significantly lower mental health and experienced more anxiety /insomnia than mothers of children without autism.

This finding was consistent with some previous research, such as Tarabek, Nouri, Yamada et al, and Salehi et al.'s study who found that mothers of children with Autism and other disabilities reported significantly less mental health than mothers of children with control group (6, 1, 10, 11).

McCarthy et al and Gray also found that mothers of children with disability experience a lot of stress (12, 13). Firat et al reported a significant increase on anxiety- tense in mothers of autism group in comparison to those of mental retarded children(5). These are similar to our finding that the mothers experience anxiety/ insomnia more than control group.

In explanation of the results it could be stated that frustration of having a disabled child, Maladaptive behaviors that characterize autism, guilt feeling resulting from anger and rejection of the child, being not aware of the causes of autism, and unknown cure, contending with the stigmatization associated with the disorder, additional expenses which can create financial burden, restrictions on

However, there aren't significant differences in other subscales- Social dysfunction, Depression, Physical symptoms. These findings are consistent with results of some studies (16, 17) which have not found differences in depression between mothers of children with special needs compared to norms or to control groups. Our results are in contrast with some other studies(1, 5, 18,19).

Perhaps this difference from the existing literature was due in part to the relatively small sample size of mothers of children with Autism in this study, or it is possible that the mothers of children with autism in the study were receiving more social support, which could decrease their complaints about social dysfunction, depression and physical symptoms. This may be due to our Iranian cultural system and support from family.

Conclusion

The diagnosis of autism affects the parent's life. The nature of the disability might affect psychological well-being of the parents. So clinicians should be aware of this issue. And don't focus on the child, only.

Limitations

Our research has some limitations; this study examined a small sample, so the

findings must be generalized with caution. Also only mothers took part in the study.

It is suggested to on larger groups of samples that will also focus on the fathers' and siblings' distress as well as those of mothers. It is also recommended to provide some programs to support families (especially mothers) that can strengthen family coping and positive adjustment.

Authors' contributions

FR and SI-M conceived and designed the evaluation, collected the clinical data, performed the statistical analysis and interpreted them. SI-M drafted the manuscript and FR revised it. Both authors read and approved the final manuscript.

References

1. Nouri M. Comparison of mental health and marital satisfaction of parents of autistic children and parents of normal children in Isfahan city. [Dissertation]: Shahid Chamran Univ.; 2009.
2. Volkmar FR, Klin A, Schultz RT, State MW. Pervasive developmental disorder. In: Sadock BG, MD and Sadock VA, MD editors. Kaplan and Sadock's Comprehensive Text Book of Psychiatry. 9th ed. Philadelphia: Lippincott Williams and Wilkins; 2009.vol2. 3540-59.
3. Gupta A, Singhal N. Psychological Support for Families of Children with Autism. Asia Pacific Disability Rehabilitation Journal (APDRJ) 2005; 16(2): 62-83.
4. Sharpley CF, Bitsika V, Efremidis B. "Influences of gender, parental health, and perceived expertise of assistance upon stress, anxiety and depression among parents of children with autism". J Intel Dev Disability 1997; 22(1): 19-28.
5. Firat S, Diler SR, Avci A, Seydaoglu G. Comparison of Psychopathology in the Mothers of Autistic and Mentally Retarded Children. J Korean Med Sc 2002; 17(5): 679-85.
6. Tarabek J. Relationship satisfaction and mental health of parents of children with autism: A comparison of autism, ADHD, and normative children, [Dissertation]: Virginia Polytechnic Institute and State Univ.; 2011.
7. Rodrigue JR, Morgan SB, Geffken, G. Families of autistic children: Psychological functioning of mothers. J Clin Child Psychol 1990; 19(4): 371- 9.
8. Noorbala AA, Bagheriyazdi SA, Mohammad K. The validation of General Health Questionnaire- 28 as a psychiatric screening tool. Hakim 2009; 11 (4):47-53.
9. Edhem D, Salem Safi P, Amiri M, Dadkhah B, Mohammadi M, Mozafari N, et al. [The examination of mental health of students in medical sciences university of Ardebil]. J Ardabil Univ Med Sci 2007; 8 (3): 229-34. Persian.
10. Yamada A, Suzuki M, Kata M, Tanaka S, Shinda T, Akechi T, et al. Emotional distress and its correlations among parents of children with pervasive developmental disorder. Psychiatry Clin Neurosci 2001; 61(6): 651-7.
11. Salehi M, Salarifard MH, Hadian M. [The compression of mental health of mothers of children with psychiatric and control group]. Iran J Psychiatry Clin Psychol 2: 78-84. Persian.
12. Mc. Carthy A, Cuskelly M, Kraayenood CE, Cohen J. Predictors of stress in mother and fathers of children with fragile x syndrome. Res Dev Disabil 2006; 27(6): 688-704.
13. Gray DE. High functioning autistic children and the construction of "normal family life". Soc Med 1997; 44(8):1097-106.
14. Mansell W, Morris KA. Survey of parents' reaction to the reaction to the diagnosis of

an autistic spectrum disorder by a local service: Access to information and use of services. *Autism* 2004; 8(4): 387-407.

15. Chimeh N, Pooretamad H, Khoramabadi R. Problems and needs of mothers with autistic children. *Fam Res* 2008; 3(3): 697-707.
16. Govender N. Attitudes of parents towards their mentally retarded children: a rural area examination. [Dissertation]: Zululand Univ.; 2002.
17. Van Riper M, Ryff C, Pridham K. Parental and family well-being in families of children with Down syndrome: a cooperative study. *Res Nurse Health* 1999; 15: 227 –35.
18. Scott BS, Atkinson L, Minton HL, Bowman T. Psychological distress of parents of infants with Down syndrome. *Am J Ment Retard* 1997; 102(2): 162-71.
19. Vesion M. Depression symptoms and emotional states in parents of disabled children. *Soc Behav Pers* 1999; 27(1): 87-97.