

## Impact of Psychological Violence on Pregnancy Outcomes in a Prospective Study

Fatemeh Abdollahi PhD<sup>\*</sup>, Farideh Rezaie Abhari MSc<sup>\*\*</sup>,  
Jamshid Yazdani Charati PhD<sup>\*\*\*</sup>, Samad Rouhani PhD<sup>\*</sup>

(Received: 1 Oct 2013; Revised: 5 May 2014; Accepted: 6 Jun 2014)

**Objective:** Violence during pregnancy has been associated with adverse pregnancy outcomes. This study aimed to explore the link between psychological violence (PSV) and pregnancy outcomes in terms of maternal and birth for the first time in women attending Mazandaran University of Medical Sciences (MUMS) primary health centers (PHCs) in Iran.

**Methods:** Prospective cohort of 1461 pregnant women exposed and non-exposed to PSV was followed until the pregnancy outcome. Modified Intimate Partner Violence, demographic and pregnancy outcomes questionnaires were administered face-to-face. Logistic regression analysis was done to estimate independent effects of the PSV on pregnancy outcomes.

**Results:** More than half of the women (69.9%) reported PSV during pregnancy. The differences between the two groups in reference with pregnancy complication did not reach statistical significance. Premature rupture of membrane was the only outcome that was independently associated with PSV.

**Conclusion:** PSV in pregnancy was frequent in our study. Although the lack of adverse pregnancy outcome following PSV was observed in this study, intervention is required to prevent the effect of violence on women and child health.

**Declaration of interest:** None.

**Citation:** Abdollahi F, Rezaie Abhari F, Yazdani Charati J, Rouhani S. Impact of psychological violence on pregnancy outcomes in a prospective study. *Iran J Psychiatry Behav Sci* 2014; 8(3): 22-7.

**Key words:** • Outcomes • Pregnancy • Psychological • Violence

### Introduction

The most shameful human rights violation is violence against women (1). Published documents have been estimated the prevalence of psychological violence (PSV) during pregnancy ranged from 19.2%, 35% to 60.5% in different parts of Iran (2-4). If these percentages were applied to the 1,365,480 pregnant women in Iran in 2011, approximately 259,441 and 819,288 women were subjected to PSV during their

pregnancies. It seems that violence might be more frequent than other medical problems during pregnancy such as preeclampsia or gestational diabetes (5).

Numerous studies have been investigated the outcomes of Intimate Partner Violence (IPV) during pregnancy, still little has been focused on PSV with positive association (6, 7). A recent study on 976 pregnant women found that the incidence of low birth weight infants was significantly increased in women who reported PSV in comparison to the non-violent group (7.6% vs. 5.1%) (8).

Early detection of PSV is important for the physical and mental well-being of mother and her child. To our knowledge, no study has been solely conducted on whether PSV against women is associated with pregnancy outcomes in Iran. This longitudinal cohort study aimed to answer two questions: a. what are the risk and protective factors for PSV against women during pregnancy, and b. what

**Authors' affiliation:** \* Department of Public Health, School of Health, Mazandaran University of Medical Sciences, Sari, Iran.  
\*\* Department of Midwifery, School of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari. \*\*\* Psychiatry and Behavioral Research Center, Addiction Institute, Department of Statistics, School of Health, Mazandaran University of Medical Sciences, Sari, Iran

• **Corresponding author:** Samad Rouhani PhD, Department of Public Health, School of Health, Mazandaran University of Medical Sciences, Sari, Iran. Address: Payambar Azam Campus, Farah Abad Rd., Sari, Iran, PO Box: 48188.  
Tele: +98 1513543748  
Fax: +98 1513542473  
Email: sr485@yahoo.com

is the role of PSV on pregnancy outcomes in terms of maternal and birth outcomes?

## Materials and Methods

### Procedures

This cohort study was conducted in the primary health centers (PHCs) of Mazandaran University of Medical Sciences (MAZUMS), the only public health centers providing prenatal care to women living in Mazandaran province in the north of Iran. Sample size was calculated based on reported prevalence of IPV in Iran and using G-power software (9). Stratified sampling method was used to selection PHCs in five parts (north, south, west, east and center) of each 16 cities. In a Poisson random method, 1500 eligible pregnant women who attended to PHCs between Februarys to September 2010 were approached. Singleton pregnant women who were not competent to give informed constant were excluded from the project. At their entry to the study, reliable socio-demographics and PSV questionnaires (Cronbach's alpha 91%) were administered face-to-face by researchers who were familiar with the project. The women were divided into two groups; who screened positive for PSV and screened negative for PSV and then women were followed-up till outcome of pregnancy. The project was approved by Mazandaran University of Medical Sciences' ethics committee.

### Psychological violence

PSV was assessed using Iranian cultural adapted of World Health Organization (WHO) Domestic Violence Questionnaire that used in Iran-Tehran before and including 14 questions on PSV (4). They consist of: (a) Are you afraid of your husband? (b) Has he threatened your life? (c) Has he threatened to hurt you or anybody close to you? (d) Does he abuse you emotionally? (e) Does he use offensive language? (f) Has he verbally abused your family with or without their presence? (g) Does he disapprove of your beliefs and principles? (h) Does he curse children? (i) Does he abandon you and your children? (j) Does he keep you from going out of the home? (k) Does he keep you from going to friends or relatives, ceremonies or other places you like to go? (l) Does he keep

you from getting a job? (m) Does he keep you from studying? (n) Does he keep you short in terms of money, food, and clothing? The PSV was considered "severe" if questions (a) and or (b) had a positive answer, PSV was considered "moderate" if at least five items from questions (a) to (n) were positive, and finally PSV was considered "mild" if fewer than five items were positive. This standard questionnaire was used after pretesting with a sample of 50 healthy pregnant women in PHCs with reliability of 0.78 Cronbach's alpha.

### Pregnancy outcomes

During the index of pregnancy, women were asked about maternal complications that included abortion (fetal loss before 20<sup>th</sup> gestation week), placenta abruption (placenta separation during pregnancy), vaginal bleeding (hemorrhage in the second or third trimester) and diabetes. Birth certificates provided information on complications of delivery and birth that contained intra uterine fetal death (IUFD) (fetal loss after 20-week of gestation), intra uterine growth retardation (IUGR; birth weight was at the 10<sup>th</sup> percentile or less based on published birth weights in weekly gestational age categories), premature rapture of membrane (PROM) (spontaneous ruptured of membrane four hours before beginning of delivery contraction), type of delivery (vaginal and sectarian section), pre-term delivery (live birth delivered before 37<sup>th</sup> week of gestations), and low birth weight (LBW) (birth weight less than 2500 gram) (10).

### Potential confounders

Potential confounders of the association between PSV during pregnancy and adverse maternal and birth outcomes were including age, years married, education and occupational status of the women and their husbands, family income, status of accommodation, parity and history of infertility. Iranian classification of income was used which classifies household income into three different categories: (low; less than 300 U.S. dollar, medium; 350-450 USD and high; more than 450 USD).

### Statistical analysis

We initially explored the prevalence

distribution of three groups of PSV (mild, moderate, severe) during pregnancy. Then, women were placed into two categories; one category entails women who were being PSV, and reference category involving women who were not. The prevalence of PSV in each level of maternal characteristics variables with its exact 95% confidence interval obtained from a binomial distribution was reported.

To investigate whether a relationship existed between the adverse pregnancy outcomes and PSV, simple logistic regression was used generating the corresponding odds ratios (OR) coupled with the 95% confidence intervals (CI). Then multiple logistic regression analysis was done to estimate independent effects of the PSV and kept in the model if P-value was 0.05 or less. These associations were adjusted for all confounders. Statistical Analysis software (SAS<sup>®</sup>), version 20.0 was used for the statistical analysis.

## Results

Data were collected through interviews with the 1461 women who were consented and followed-up the study. A small number of women (n = 39) refused to participate. Among the participants, 1,020 (69.9%) reported that they were violated psychologically by their husband during the pregnancy. The most common PSV was kept them short in terms of money, food, and clothing (88.8%).

Distribution of three groups of PSV (mild, moderate, and severe) during pregnancy is shown in table 1. More women fell in mild group of PSV (39.9%). The results of unadjusted analysis of the factors that affect on PSV are illustrated in table 2.

**Table 1.** Prevalence severity of psychological violence during pregnancy

Violence	Number (%)
No	439 (30.1)
Mild	582 (39.9)
Moderate	107 (7.3)
Severe	331 (22.7)

**Table 2.** Prevalence of psychological violence (PSV) during pregnancy by maternal characteristics

Characteristics <sup>†</sup>	Violence (n = 439) Number (%)	Non-violence (n = 1,020) Number (%)	OR (95% CI) <sup>‡</sup>
<b>Age (years)</b>			
< 25	460 (46.7)	204 (46.3)	1.01 (0.81-1.27)
≥ 25	525 (53.3)	237 (53.7)	
<b>Gravida</b>			
Primiparous	23 (2.3)	14 (3.2)	0.72 (0.37-1.42)
Multipara	962 (97.7)	426 (96.8)	
<b>Educational status</b>			
Illiterate and primary	344 (34.9)	167 (37.9)	0.88 (0.69-1.11)
Secondary and over	641 (65.1)	274 (62.1)	
<b>Occupational status</b>			
Housekeeper	815 (84.9)	372 (85.1)	0.93 (0.67-1.29)
Employed	145 (15.1)	62 (14.3)	
<b>Age of husband</b>			
<30	546 (55.4)	247 (56)	0.97 (0.77-1.22)
≥30	439 (44.6)	194 (44)	
<b>Husband's educational status</b>			
Illiterate and primary	369 (37.5)	152 (34.5)	1.13 (0.9-1.43)
Secondary and over	615 (62.5)	288 (65.5)	
<b>Husband occupational status</b>			
Governmental job	805 (81.7)	372 (84.5)	0.81 (0.6-1.10)
Non- governmental job	180 (18.3)	68 (15.5)	
<b>Polygamy</b>			
Yes	59 (6)	33 (7.5)	1.26 (0.81-1.97)
No	926 (94)	408 (92.5)	
<b>Duration of marriage (years)</b>			
< 5	611 (62)	265 (60.2)	1.07 (0.85-1.35)
≥ 5	374 (38)	175 (39.8)	
<b>Having relative kinship with husband</b>			
Yes	133 (13.6)	73 (16.7)	0.78 (0.57-1.06)
No	848 (86.4)	363 (83.3)	
<b>Housing</b>			
Own house	412 (41.8)	202 (45.8)	1.17 (0.93-1.47)
Renting	573 (58.2)	239 (54.2)	
<b>Total household income (monthly)</b>			
High (more than 450 USD; U.S. dollar)	223 (22.7)	97 (22)	1.03 (79-1.36)
Low (350 USD and less)	760 (77.3)	343 (78)	

<sup>†</sup> There was no significant difference between two groups; <sup>‡</sup> Result from chi-square test

The characteristics of women were same in violated and nonviolent groups. Table 3 shows the association between PSV during index of pregnancy and pregnancy outcomes after adjusting. PSV had no statistically significant impact on pregnancy outcomes except PROM. There was a statistical significant difference between the violated and nonviolent women in terms of PROM [OR: 0.65 (CI: 0.43-0.97)].

**Table 3.** Association between pregnancy outcomes and psychological violence during pregnancy (adjusted analysis; n = 1,469)

Outcome	OR (95% CI)	P-value
Abortion	0.93 (0.55-1.55)	0.78
Diabetes	1.41 (0.63-3.15)	0.40
BP <sup>†</sup>	3.6 (0.44-28.8)	0.22
Hemorrhage	1.12 (0.49-2.56)	0.78
Placenta abruption	0.62 (0.27-1.41)	0.25
Preeclampsia	0.99 (0.59-1.64)	0.97
Term vs. pre-term	0.84 (0.61-1.17)	0.32
Vaginal delivery vs. C/S <sup>‡</sup>	1.25 (0.97-1.61)	0.08
PROM <sup>§</sup>	0.65 (0.43-0.97)	0.03
Low Birth Weight	0.71 (0.5-1.01)	0.059

<sup>†</sup> Blood pressure; <sup>‡</sup> Cesarean delivery

<sup>§</sup> Premature rupture of membrane

## Discussion

To our knowledge, this is the first cohort study to investigate the impact of PSV solely on pregnancy outcomes in Iran. It is not easy to measure PSV because of cultural sanctioning, reluctant women to reveal it and no standardized definition of PSV (11, 12).

We found that PSV was common (69.9%) among pregnant women attending PHCs. This is comparable to the findings of previous studies conducted in Iran and indicated that PSV is more common than physical violence (60.5% and 57% vs. 14.6% and 5.5%) (4, 13). A population-based study in Hong Kong found around three-fourths of the violence during pregnancy (216 of 296 violated women) was psychological only (7). It is possible that the psychological violence could be a forerunner of other forms of violence. During pregnancy staying in touch frequently with the health service professionals provides an appropriate chance to detect who are at risk. Given the offer that progression from PSV to other forms of violence takes a short time, identifying those at risk and carry out the relevant interventions is noticeable. Further studies are needed to determine if

violence is predominantly psychological in nature especially in the traditional nations.

The lack of adverse pregnancy outcome following PSV was observed in our study. The only similar findings were observed in a cross-sectional study in Hong Kong by Tiwari et al. (7). The authors found no association between PSV and LBW, pre-term birth, type of delivery, low Apgar score and admission the neonate to the hospital.

The research findings on the potential association between PSV solely and pregnancy outcomes are rare and inconclusive. Many researches explored this association in combination of all types of violence (physical, psychological and/or sexual). Zareen et al. found no statistical association between all types of violence and maternal and perinatal complications (14). This was in consistent with the findings of Covington et al., where no increase in LBW and pre-term birth was found in women who subjected to violence (15).

Certainly, the experience of violence during pregnancy has many adverse outcomes including LBW, preterm birth and premature rupture of membrane (PROM) (2, 13, 16, 17). In agreement with the above reports, positive association was found between violence and PROM in the present study. Many factors may explain this association including direct physical and health effects with mental health factors as well (12). Adverse effects are not restricted to women who subject only physical violence, as even PSV has been associated to difficult pregnancy outcomes. Women who are subjected to violence during pregnancy may experience higher rate of sexually transmitted disease (STDs) (12) that expose them at increased risk for PROM (10).

Although the present study found the lack of adverse effect of PSV on pregnancy outcomes, the psychological anger between partners can have negative effect on all family members in particular on infants (7). Thus, screening for PSV during pregnancy is needed and intervention is considered necessary not only to protect the women from adverse effects of violence, but also reduce the harm on the children. Moreover, health care providers could be able to provide

limited counseling to motivational interviewing or increasing women's self-esteem and a sense of inner control. More comprehensive research is needed to more completely understand the IPV problem and to find appropriate solutions.

As the women participated voluntarily in this study, selection bias may have occurred. The other limitation of the study is that an episode of violence prior to the pregnancy and following the interview may have been missed. Furthermore, this study failed to provide support to the women who were subjected to PSV.

### Acknowledgments

This study was supported by MUMS in Iran. We are thankful to healthcare providers and mothers who participated in this project.

### Authors' contributions

FA designed the study, re-evaluated the data, advised on the analysis and drafted the manuscript. SR, FRA and JYCH advised on the study design, helped to collect the data, analysis and interpretation of the data and revised the manuscript. All authors read and approved the version submitted.

### References

- Garcia-Moreno C, Jansen HA, Ellsberg M, Heise L, Heise L. WHO multi-country study on women's health and domestic violence against women. Geneva, Switzerland: World Health Organization; 2005.
- Faramarzi M, Esmaeilzadeh S, Mosavi S. Prevalence, maternal complications and birth outcome of physical, sexual and emotional domestic violence during pregnancy. *Acta Medica Iranica* 2005; 43(2): 115-22.
- Salari Z, Nakhaee N. Identifying types of domestic violence and its associated risk factors in a pregnant population in Kerman hospitals, Iran Republic. *Asia Pac J Public Health* 2008; 20(1): 49-55.
- Jahanfar Sh, Malekzadegan Z. The prevalence of domestic violence among pregnant women who were attended in Iran University of Medical Science Hospitals. *J Fam Violence* 2007; 22(8): 643-8.
- Gazmararian JA, Lazorick S, Spitz AM, Ballard TJ, Saltzman LE, Marks JS. Prevalence of violence against pregnant women. *JAMA* 1996; 275(24): 1915-20.
- Street AE, Arias I. Psychological abuse and posttraumatic stress disorder in battered women: examining the roles of shame and guilt. *Violence Vict* 2001; 16(1): 65-78.
- Tiwari A, Chan KL, Fong D, Leung WC, Brownridge DA, Lam H, et al. The impact of psychological abuse by an intimate partner on the mental health of pregnant women. *BJOG* 2008; 115(3): 377-84.
- Yost NP, Bloom SL, McIntire DD, Leveno KJ. A prospective observational study of domestic violence during pregnancy. *Obstet Gynecol* 2005; 106(1): 61-5.
- Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power analyses using G\*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009; 41(4): 1149-60.
- Cunningham F, Leveno K, Bloom S, Hauth J, Gilstrap L, Wenstrom K. Williams obstetrics. 22<sup>th</sup> ed. Philadelphia, PA: McGraw-Hill Professional; 2005.
- Nasir K, Hyder AA. Violence against pregnant women in developing countries: review of evidence. *Eur J Public Health* 2003; 13(2): 105-7.
- Bailey BA. Partner violence during pregnancy: prevalence, effects, screening, and management. *Int J Womens Health* 2010; 2: 183-97.
- Khosravi F, Hashemi Nasab L, Abdollahi M. [Study of the incidence of domestic violence among pregnant women referring to childbirth unit of Sanandaj Hospitals.] *Urmia Med J* 2008; 19(1): 8-14. Persian
- Zareen N, Majid N, Naqvi S, Saboohi S, Fatima H. Effect of domestic violence on pregnancy outcome. *J Coll Physicians Surg Pak* 2009; 19(5): 291-6.
- Covington DL, Justason BJ, Wright LN. Severity, manifestations, and consequences of violence among pregnant

- adolescents. *J Adolesc Health* 2001; 28(1): 55-61.
16. Ferri CP, Mitsuhiro SS, Barros MC, Chalem E, Guinsburg R, Patel V, et al. The impact of maternal experience of violence and common mental disorders on neonatal outcomes: a survey of adolescent mothers in Sao Paulo, Brazil. *BMC Public Health* 2007; 7: 209.
17. Shah AJ, Kilcline BA. Trauma in pregnancy. *Emerg Med Clin North Am* 2003; 21(3): 615-29.