

## An Atypical Suicide Attempt: Self-Inflicted Intra-Cardiac Injury with Sewing Needle

Nazila Shahmansouri MD<sup>\*</sup>, Mahmood Shirzad MD<sup>\*\*</sup>, Sam Zeraatianejad Davani MD<sup>\*\*\*</sup>,  
Fetemeh Heidari MA<sup>\*\*\*\*</sup>

(Received: 5 Junm 2014; Revised: 13 Oct 2014; Accepted: 24 Dwc 2014)

Atypical cases of suicide are less likely to be seen in general hospitals, nonetheless require further investigation into the precipitating factors as well as proper follow-up. This paper illustrates a 61-year-old woman with major depressive disorder who experienced auditory hallucinations during delirious periods of taking low-dose benzodiazepines, who referred to the hospital with a sewing needle stuck in her chest wall. The needle was successfully removed. Psychiatric problems are often underdiagnosed, therefore undertreated in general hospitals. Thus close monitoring of the patients during the hospitalization and after discharge is crucial.

**Declaration of interest:** None.

**Citation:** Shahmansouri N, Shirzad M, Zeraatianejad Davani S, Heidari F. An atypical suicide attempt: self-inflicted intra-cardiac injury with sewing needle. *Iran J Psychiatry Behav Sci* 2014; 8(4): 80-2.

**Keywords:** Atypical Suicide • Self-Inflicted Injury • Surgery

### Introduction

The two most common suicide methods are self-poisoning and self-injury, respectively (1). Suicidal attempt is a risk factor for successful suicide (2). Studies have shown that 24-42% of those who attempted suicide had planned in less than 5 minutes (2). In some studies, young age and history of physical illness in females have been demonstrated as suicide risk factors (3). Many suicide victims suffer from physical illness and psychiatric disorders (2).

Psychiatric problems are often underdiagnosed, and therefore, undertreated in general hospitals (4, 5). Moreover, unconventional suicide methods can increase diagnostic and therapeutic complications. This paper reports a rare suicide method

(penetrating sharp foreign bodies into the heart) of which only a few similar cases were found in the literature (6, 7). Most cases of self-harm are demonstrations of underlying psychiatric problems (8).

### Case Report

A. P., a 61-year-old woman, was taken to the emergency ward of the Tehran Heart Center, Iran, by her husband and son. On admission, she was suffering from dyspnea, and there was a sewing needle penetrating her chest. During the physical examination, the heart and lungs were normal. Emergency fluoroscopy, echocardiography and computed tomography (CT) scan were performed. The chest CT scan revealed that the needle had penetrated into the heart, and echocardiography demonstrated that the needle had ruptured the pericardium, but it had not entered the left ventricle. The patient underwent surgery 4 hours after admission, and the needle was removed successfully.

An interview with her family disclosed a history of major depressive disorder and pharmacological treatment. During the first 3 sessions of psychiatric counseling after the surgical operation, the patient denied a suicide

**Authors' affiliation:** \* Assistant Professor, Department of Psychiatry, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran. \*\* Assistant Professor, Department of Cardiovascular Surgery, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran. \*\*\*Cardiac Surgery Fellowship, Department of Cardiovascular Surgery, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran. \*\*\*\*Psychologist, Tehran Heart Center, Research Branch, Tehran, Iran.

• **Corresponding author:** Nazila Shahmansouri, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran  
Tel: +98 2188029600  
Fax: +98 2188029731  
Email: n.shahmansouri@gmail.com

attempt. She claimed that an accidental fall was the cause of her injury. On the 4<sup>th</sup> session, however, she admitted that the problem was not accidental, stating that she had attempted unplanned impulsive suicide by swallowing needles twice before, once 10 years and once a year previous to this. In both cases, she had been hospitalized for some days and discharged without any complications.

The patient had been suffering from major depressive disorder since the sudden death of her son, 10 years ago. During the previous 5 years, she had received different pharmacological treatments, namely clonazepam, alprazolam, venlafaxine, and Depakine, which were not followed through. No history of psychotherapy or electroconvulsive therapy was reported.

The patient reported having experienced auditory hallucinations during different periods of taking low-dose benzodiazepine pills. According to her family, during these periods, the patient had experienced disorientation, disturbed consciousness, distraction, and auditory hallucinations. Through these accounts we were guided toward the delirious picture of those episodes in her past. She also self-medicated with benzodiazepine pills in those periods reported by the family, as she was not receiving adequate attention from them.

The family members described her as a demanding and irritable person who needs a great deal of attention, and is also prone to impulsive behavior when under stress or not receiving adequate attention.

After thorough examination, venlafaxine was prescribed as antidepressant. The patient should have remained hospitalized until the completion of the treatment, but the patient as well as her family members consented to leave the hospital after 5 days despite the advice of the medical team. Efforts to convince her to adhere to the course of treatment failed due to her refusal to do so.

### Discussion

Violent methods of suicide attempts are uncommon. In this case, the patient initially denied having attempted suicide, but finally

disclosed her previous suicide attempts and depressive symptoms.

Our patient's bizarre attempt at taking her own life could be explained in the following two ways:

1. Her personality traits of being demanding, impulsive, and sensitive to self-perceived inadequate attention might have rendered her prone to these attempts. This is evidenced by her impulsive and self-prescribed consumption of benzodiazepine pills as a result of feelings ignored as reported by her family and significant other. Usually, individuals with higher impulsivity are at a greater risk of attempting suicide (9).

2. Swallowing needles or sticking them into the chest wall could be an instance of self-harm inasmuch as an impulsive person is liable to exhibit self-harming behavior (9). Nevertheless, the possibility of suicide attempts should not be discounted.

Unfortunately, our patient's refusal to permit follow-up prevented us from further looking into this unusual case. This study highlights the necessity of a multi-disciplinary approach to examining suicide cases. It is advisable to consider surgical cases which appear unusual to the surgeon as possible instances of self-harm or suicide attempts and to place them under close clinical observation. In general hospitals, the psychiatric background of the patient is likely to be overlooked, and as a result, suitable treatment is not provided (5). In most regions of the world, less than half of the suicide cases are estimated to receive the necessary medical attention (10). Alarming, this is inconsistent with the finding that these individuals tend to have further self-harming attempts in their lives (11). Patients with no previous suicidal ideation, depressive mood, or hopelessness, no history of substance or alcohol abuse, and strong social network and support are at lower risk of attempting suicide (12). Close monitoring of patients during hospitalization is crucial. The majority of successful suicides occur shortly after hospital discharge (13). Ignoring patients' psychiatric history and impulsive suicidal attempt after discharge lead to the underestimation of the suicide risk

of patients admitted to general hospitals (13).

### Acknowledgments

The authors would like to thank staff of ICU open heart of the Tehran Heart Center.

### Authors' contributions

NSH and FH wrote the manuscript. MSH and SZD carried out the cardiologic studies and surgery. NSH and FH have been involved in the acquisition of clinical data/psychiatric evaluation and reviewing the scientific literature. All authors revised the manuscript critically for important intellectual content. NSH contributed to the final version and carried out the clinical case report. All authors read and approved the final manuscript.

### References

1. Lin CJ, Lu HC, Sun FJ, Fang CK, Wu SI, Liu SI. The characteristics, management, and aftercare of patients with suicide attempts who attended the emergency department of a general hospital in northern Taiwan. *J Chin Med Assoc* 2014; 77(6): 317-24.
2. Wei S, Liu L, Bi B, Li H, Hou J, Chen W, et al. Comparison of impulsive and nonimpulsive suicide attempt patients treated in the emergency departments of four general hospitals in Shenyang, China. *Gen Hosp Psychiatry* 2013; 35(2): 186-91.
3. Wei S, Yan H, Chen W, Liu L, Bi B, Li H, et al. Gender-specific differences among patients treated for suicide attempts in the emergency departments of four general hospitals in Shenyang, China. *Gen Hosp Psychiatry* 2013; 35(1): 54-8.
4. Salehian O, Teoh K, Mulji A. Blunt and penetrating cardiac trauma: a review. *Can J Cardiol* 2003; 19(9): 1054-9.
5. Kadis P, Pogorevc L, Sipek M, Vidovic D. Unusual attempted suicide by shooting through heart. *Forensic Sci Int* 2005; 147(Suppl): S29-32.
6. DuVal CJ. Improving nursing satisfaction. *Nurs Manage* 1992; 23(6): 64.
7. Chui WH, Cheung DL, Chiu SW, Lee WT, He GW. A non-fatal impalement injury of the thorax. *J R Coll Surg Edinb* 1998; 43(6): 419-21.
8. Schechter DC, Gilbert L. Injuries of the heart and great vessels due to pins and needles. *Thorax* 1969; 24(2): 246-53.
9. Maser JD, Akiskal HS, Schettler P, Scheftner W, Mueller T, Endicott J, et al. Can temperament identify affectively ill patients who engage in lethal or near-lethal suicidal behavior? A 14-year prospective study. *Suicide Life Threat Behav* 2002; 32(1): 10-32.
10. Bertolote JM, Fleischmann A, De LD, Bolhari J, Botega N, De SD, et al. Suicide attempts, plans, and ideation in culturally diverse sites: the WHO SUPRE-MISS community survey. *Psychol Med* 2005; 35(10): 1457-65.
11. Bille-Brahe U, Jessen G. Repeated suicidal behavior: a two-year follow-up. *Crisis* 1994; 15(2): 77-82.
12. Ronquillo L, Minassian A, Vilke GM, Wilson MP. Literature-based recommendations for suicide assessment in the emergency department: a review. *J Emerg Med* 2012; 43(5): 836-42.
13. Cheng IC, Hu FC, Tseng MC. Inpatient suicide in a general hospital. *Gen Hosp Psychiatry* 2009; 31(2): 110-5.