

## Cigarette Smoking among Iranian University Students: Reasons and Attitudes

Manzume Shamsi Meymandi, Pharm. D<sup>\*</sup>, Nouzar Nakhac, MD<sup>\*\*</sup>  
Kouros Divsalar, MSc<sup>\*</sup>, Gioia Heravi, MD<sup>\*\*</sup>

(Received: 27 Dec 2009 ; Accepted: 7 Aug 2010)

**Objective:** According to the fact that university students are role models for the whole society, this study was performed to evaluate reasons of and attitudes toward cigarette smoking among Iranian university students.

**Methods:** This study was performed as a cross-sectional self-reported questionnaire based study among university students of Kerman city of Iran. Our questionnaire was attributed to 1750 university students who were selected randomly. Current smoking was defined as using cigarette within the last 30 days. Participants' demographic data in addition to their reasons for and attitudes toward smoking were obtained. Frequency, distribution and factor analysis of attitude items followed by a varimax rotation was used to evaluate the attitudinal statements.

**Results:** From 1750 university students, who filled our questionnaire, 31% were ever smokers (51% males and 15% females) and 11% were current smokers (21.5% males and 2% females). Avoiding depression and taking pleasure were the reasons mentioned as important reasons the most frequently. Besides, the most frequently attitudinal statements were as follows: "My parents upset knowing my habit" and "Cigarette causes disease". Furthermore, the mean score of encouraging attitudes was significantly higher among males, current smokers and those with a tendency toward cigarette smoking.

**Conclusion:** This study confirmed that addressing attitudes toward smoking while developing preventive strategies against it among university students is an obvious necessity.

**Declaration of interest:** None.

*Iranian Journal of Psychiatry and Behavioral Sciences (IJPBS), Volume 4, Number 2, Autumn and Winter 2010: 37-41.*

**Keywords:** Attitude • Cigarette Smoking • University Student

### Introduction

Tobacco use is one of the major preventable causes of death in the world and is one of the most prevalent public health problems in Iran. The rate of cigarette smoking among Iranian adults with 19 to 49 years of age was reported to be 21% in men and 7% in women (1). On the other hand, prevalence of cigarette smoking is shown to have an increasing trend in developing countries in recent years (2,3). According to role- modeling of university students for children and adolescents, their tobacco use may increase the cigarette consumption in whole

society consequently (4,5). Indeed, cigarette smoking of university students can even be considered as a useful index of tobacco use among adults (5).

In spite of different studies which investigated the prevalence and risk factors of smoking among youths in Iran (1,3,6-10), limited information is available regarding this issue in Iranian university students compared with other developing countries (11). Thus, underlining the role modeling of university students, this study intends to evaluate reasons of and attitudes toward cigarette smoking among Iranian university students.

### Materials and Methods

In this cross- sectional study, which was performed in 2006, data were gathered from 1750 randomly selected students of two universities of Kerman. Kerman city is located

**Authors' affiliations :** \*\* Kerman University of Medical Sciences, \* Kerman Neuroscience Research Center,

•**Corresponding author : Manzume Shamsi Meymandi, MD,**  
Kerman Neuroscience Research Center, Jahad Blvd., EbneSina St.,  
Kerman, Iran  
Tel : +98 341 2264180  
Fax : +98 341 3221671  
E-mail: manzume@yahoo.com

in the south west of Iran, 1000 kilometer far from the capital, and has a population of more than half million. This study was approved by ethical research committee of Kerman University of medical sciences.

A self- reported questionnaire including questions on demographic data such as age, gender and marital status, six questions on frequency of cigarette smoking, seven questions on reasons and ten questions on attitudes toward smoking was used in the present study. Current smoking was defined as cigarette use during the last 30 days (12).

The questionnaire was distributed to volunteer participants by research personnel during university classes on general courses, and after a brief explanation of aim and scope of the present study. Anonymous filled in questionnaires were gathered in the same section. The content validity of our questionnaire was confirmed based on literature review and an expert panel. For evaluating its reliability the test- retest method was used in a pilot study where interclass correlation coefficient was less than 0.8.

Data analysis was carried out using Statistical package for social sciences (SPSS) software, version 17. Statistical analysis included frequency, distribution and factor analysis of attitude items. Principal axis factor analysis followed by a varimax rotation was used to identify key dimensions of the attitudinal statements. T-test was then used to compare the mean score of attitudes between the two groups of smoker and non-smoker university students.

## Results

From 1750 students of Kerman University who were invited to participate in this study, 1719 students completed the questionnaires, with a response rate of 98%. Forty two of these filled in questionnaires were excluded due to their incomplete data and 1677 questionnaires were finally enrolled.

The mean age of participants was  $21.2 \pm 2.1$  years, most of whom were single (92%) and females constituted 52% of participants. Five hundred and nineteen of participants (41%) were intended to smoke, while 516 of them

(31%) were experienced cigarette smoking during their lifetime. A total number of 184 (11%) were current smokers, from whom 88 percent were male. Approximately 2% of current smokers consumed more than 10 cigarettes per day.

The frequency of agreement with each of attitudinal statements is shown in Table 1. More than 90% of participants believed that their parents would be upset knowing cigarette smoking habit of them, and believed that cigarette smoking causes diseases.

**Table 1:** University students' attitudes toward cigarette smoking

| Phrases  | Agree       | Disagree    |
|--|-------------|-------------|
| Cigarette smoking is a normal habit                | 26%(445)    | 74%(1205)   |
| I accept a cigarette from my close friend          | 11.8%(206)  | 88.2%(1444) |
| Cigarette smoking develops social personality      | 3.6%(63)    | 96.4%(1684) |
| I tolerate who smoke near me                       | 38%(665)    | 62%(985)    |
| Cigarette smoking is expensive                     | 73.2%(1281) | 26.8%(469)  |
| Cigarette smoking causes diseases                  | 92.6%(1620) | 7.4%(130)   |
| Even smoking one cigarette might lead to addiction | 44.2%(773)  | 55.8%(977)  |
| Cigarette smoking destroys the beauty              | 55.3%(968)  | 44.7%(782)  |
| My parents upset knowing my smoking habit          | 94.7%(1657) | 5.3%(93)    |

"Encouraging" attitudes are in italic while "Discouraging" attitudes are in bold.

On the other hand, avoiding depression (42.4%) and taking pleasure (41.8%) were considered as the most important reasons of smoking by university students (Table 2).

**Table 2:** The frequency of importance of reasons toward smoking in university students

| Reasons                | Important  |
|------------------------|------------|
| Avoiding depression    | 42.4%(742) |
| Taking pleasure        | 41.8%(732) |
| To relief stress       | 13.6%(238) |
| To relief anger        | 0.5(9)     |
| Easier problem solving | 0          |

Principal axis factor analysis of attitude items followed by a varimax rotation indicated that attitudes could be categorized into two types. One category which includes positive attitudes is named as encouraging, whereas another category containing negative attitudes is named discouraging. The Cronbach's alpha for both dimensions was approximately 0.60. The mean score of encouraging attitude was significantly higher in males, current smokers, ever smokers and those with a tendency toward cigarette smoking compared with those who have discouraging ones (Table 3).

**Table 3:** Mean of encouraging and discouraging attitude scores in different group of university students

|                                  | Discouraging Scores | Encouraging Scores |
|----------------------------------|---------------------|--------------------|
| Non Current smokers              | 3.13 ± 0.01         | 1.55 ± 0.01        |
| Current smokers                  | 2.66 ± 0.04         | 2.42 ± 0.04*       |
| Male                             | 2.93 ± 0.02         | 1.81 ± 0.02*       |
| Female                           | 3.19 ± 0.02         | 1.52 ± 0.02        |
| Tendency to cigarette smoking    | 2.88 ± 0.04         | 1.92 ± 0.02*       |
| No Tendency to cigarette smoking | 3.21 ± 0.01         | 1.46 ± 0.01        |
| Ever smoker                      | 2.81 ± 0.02         | 2.01 ± 0.03*       |
| Never smoker                     | 3.20 ± 0.01         | 1.50 ± 0.01        |

Data are expressed as Mean ± SE of scores.

\*p<0.001 indicates a statistically significant difference between discouraging and encouraging attitudes.

## Discussion

This study was performed to assess the reasons of and attitudes toward cigarette smoking among university students of Kerman city of Iran. Confidentiality and autonomy of participants were respected to ensure the validity of data.

One hundred and eighty four (11%) of 1677 enrolled questionnaires were filled by current smokers, with a mean age of  $21.2 \pm 2.1$  years. Besides, the mean age of first smoking experience which was shown to be  $14.5 \pm 2.4$  and  $13.2 \pm 2.5$  years among Iranian adolescents in 2006 and 2007 respectively has been confirmed in this study (13,14). On the other hand, this variable was shown to be  $17.94 \pm 1.64$  in another study among Iranian university students in 2008 (6). The mean age of smoking initiation is reported to be 19 years in Syria, 17 years in Brazil and 14 years in USA (15,16).

As mentioned before, prevalence of smoking was reported to be approximately 11% in the present study. This result is the same as the prevalence of smokeless tobacco use among students in Belucestan, which is a neighbor province of Kerman (6). This observation is in accordance to the obtained data of Iran's general population (1). In contrast, this prevalence was markedly fewer than the results of similar studies among university students of Syria (18.6%), Turkey (49.4%), and USA (28.5%) (15-17).

Moreover, a relation was found between cigarette smoking and gender in the present study (1,13,18). Similar results have been obtained

in previous studies in Iran and its neighbors. It means that male students were more likely to be life time smokers than females (3,6). Controversially, in an international study conducted in 23 countries, the prevalence of current smoking in males (34%) was similar to females (27%). This prevalence was even shown to be the same in both genders in some studies (5).

Most of participants declared that smoking makes smokers ugly, leads to addiction and causes diseases. These statements were similar to Karachi students' beliefs (18). It was encouraging to see that students thought that smoking was injurious to health. Indeed, it means that students' knowledge of smoking's side effects and the probability of its dependency was high (table 1). From the other side, 38% of participants tolerated breathing in someone else's secondhand smoke and 26% considered smoking as a normal habit. It can be claimed that in spite of good level of students' knowledge, a social tolerance with positive attitudes toward smoking still exists. This positive attitude was seen in another study conducted among adolescents in Iran (8).

According to the fact that frequency and development of cigarette smoking is significantly predictable through attitudes (19), it is important to understand the factors that might influence students' smoking behaviors. Regarding our findings increasing discouraging factors along with decreasing encouraging factors might change subsequent behaviors of university students in future, while increasing the cost of cigarette would not influence their future smoking behaviors. As most of university students in Iran are financially dependent on their parents, the cost of cigarettes could be only a limiting factor until they become independent and find a job in future. Increasing students' level of knowledge would not be effective either because the number of smoker students increases during studying years at university (20). Moreover, people with higher level of education have shown to smoke more (7). It is paradoxical that students begin and continue cigarette smoking in spite of their increased knowledge of its risk factors (20). This may reflect that the important determinant for smoking is the social acceptance of

smoking. Hence, social personality was one of our encouraging factors on which in dept studies should be performed.

Parental pressure was another important item that can discourage students from smoking habit. This might also lead to adolescents' hiding of tobacco use (3). There is a significant positive correlation between parental beliefs and risk of smoking among students (21). From the other side, a significant association has been found between students' having a smoker family member or a smoker friend with prevalence of smoking in them (3). Furthermore, although parental smoking might influence the initial stage of smoking attempts, the following social conformation is more dependent on smoking habits of peers, siblings and best friends (22,23).

University students' attitudes toward encouraging and discouraging factors are summarized in Table 1. Personal characteristics such as self presentation and social behaviors in addition to their beliefs were confirmed as smoking initiative risk factors in a previously performed qualitative study (24). In this regard, "having social personality" and considering smoking as an acceptable and tolerable behavior, especially when it was offered from a "close friend", leads to exclusive group formation. Peer group identity has been mentioned as an important developmental issue previously. The influence of role modeling on development of personal traits and formation of personal habits was cited in previous studies as well (14,25). Additionally, "strain theory" might influence cigarette smoking culturally (26). Hence, we suggest more social skills training approaches rather than providing university students with more information.

In conclusion, university period appeared to have critical influences on the development of social personality and identity formation. A strong association has been confirmed between youths' smoking and adult's substance use which impacts inevitably on future social health (27). Hence, some investigators suggested school based prevention programs (28) or limitation of direct and indirect tobacco use promotion through introducing tough rules against smoking in public areas (12,18,28,29).

However, we suggest more social skills training approaches rather than providing information about harmful effects of cigarette smoking or forbidding it.

### Acknowledgment

This work is a part of a study aimed at gaining a better understanding of cigarette smoking among Iranian university students, supported by a grant (86/143) from research deputy of Kerman University of Medical sciences.

### References

1. Sarraf-Zadegan N, Boshtam M, Shahrokhi S, Naderi GA, Asgary S, Shahparian M, et al. Tobacco use among Iranian men, women and adolescents. *Eur J Public Health* 2004; 14(1): 76-8.
2. Weiss JW, Palmer PH, Chou CP, Mouttapa M, Johnson CA. Association between psychological factors and adolescent smoking in seven cities in China. *Int J Behav Med* 2008; 15(2): 149-56.
3. Eftekhari Ardebili M, Nassr M, Rassulian M, Ghalebandi M, Daneshamuz B, Salehi M. Prevalence of cigarette smoking in Tehran: a household study. *Iran J Psych Behav Sci* 2007; 1(2): 33-7.
4. Rigotti NA, Moran SE, Wechsler H. US college students' exposure to tobacco promotions: prevalence and association with tobacco use. *Am J Public Health* 2005; 95(1): 138-44.
5. Rigotti NA, Lee JE, Wechsler H. US college students' use of tobacco products: results of a national survey. *Jama* 2000; 284(6): 699-705.
6. Jalilvand M, Nikmanesh Z, Kazemi T, Emamhadi M. Smokeless tobacco use among university students: A cross-sectional study in Iran, Sistan Baloochestan province, 2008. *Iran J Psych Behav Sci* 2010; 4(1): 23-9.
7. Fotouhi A, Khabazkhoob M, Hashemi H, Mohammad K. The prevalence of cigarette smoking in residents of Tehran. *Arch Iran Med* 2009; 12(4): 358-64.

8. Mohammadpoor A, Fakhari A, Pourafkary N. Cigarette Smoking among Iranian Adolescents. *Iran J Psych Behav Sci* 2007; 1(1): 30-5.
9. Zarghami M, Khalilian AR, Veryani AR. Cigarette smoking among Sari high school students. *Med J Iran Hos* 2003; 6(1): 40-6.
10. Siam Sh. [KAP study of cigarette smoking in male high school students in Rasht.] *Behdashte Jahan* 1996; 48(3): 20-5. Persian.
11. Smith DR, Leggat PA. An international review of tobacco smoking among medical students. *J Postgrad Med* 2007; 53(1): 55-62.
12. WHO guidelines for controlling the tobacco epidemic. Geneva: WHO; 1998.
13. Kelishadi R, Ardalan G, Gheiratmand R, Majdzadeh R, Delavari A, Heshmat R, et al. Smoking behavior and its influencing factors in a national-representative sample of Iranian adolescents: CASPIAN study. *Prev Med* 2006; 42(6): 423-6.
14. Kelishadi R, Mokhtari MR, Tavasoli AA, Khosravi A, Ahangar-Nazari I, Sabet B, et al. Determinants of tobacco use among youths in Isfahan, Iran. *Int J Public Health* 2007; 52(3): 173-9.
15. Maziak W, Hammal F, Rastam S, Asfar T, Eissenberg T, Bachir ME, et al. Characteristics of cigarette smoking and quitting among university students in Syria. *Prev Med* 2004; 39(2): 330-6.
16. Steptoe A, Wardle J, Cui W, Baban A, Glass K, Tsuda A, et al. An international comparison of tobacco smoking, beliefs and risk awareness in university students from 23 countries. *Addiction* 2002; 97(12): 1561-71.
17. Oksuz E, Mutlu ET, Malhan S. Characteristics of daily and occasional smoking among youths. *Public Health* 2007; 121(5): 349-56.
18. Khan FM, Husain SJ, Laeeq A, Awais A, Hussain SF, Khan JA. Smoking prevalence, knowledge and attitudes among medical students in Karachi, Pakistan. *East Mediterr Health J* 2005; 11(5-6): 952-8.
19. Andrews JA, Duncan SC. The effect of attitude on the development of adolescent cigarette use. *J Subst Abuse* 1998; 10(1): 1-7.
20. Dumitrescu AL. Tobacco and alcohol use among Romanian dental and medical students: a cross-sectional questionnaire survey. *Oral Health Prev Dent* 2007; 5(4): 279-84.
21. Rozi S, Akhtar S, Ali S, Khan J. Prevalence and factors associated with current smoking among high school adolescents in Karachi, Pakistan. *Southeast Asian J Trop Med Public Health* 2005; 36(2): 498-504.
22. Nakhaee N, Divsalar K, Bahreini S. Prevalence of and factors associated with cigarette smoking among university students: a study from Iran. *Asia Pac J Public Health*. In Press 2009.
23. Xiang H, Wang Z, Stallones L, Yu S, Gimbel HW, Yang P. Cigarette smoking among medical college students in Wuhan, People's Republic of China. *Prev Med* 1999; 29(3): 210-5.
24. Niknami S, Akbari M, Ahmadi F, Babae-Rouchi G, Heidarnia A. Smoking initiation among Iranian adolescents: a qualitative study. *East Mediterr Health J* 2008; 14(6): 1290-300.
25. Asfar T, Ward KD, Eissenberg T, Maziak W. Comparison of patterns of use, beliefs, and attitudes related to waterpipe between beginning and established smokers. *BMC Public Health* 2005; 5: 19.
26. Carvajal SC, Wiatrek DE, Evans RI, Knee CR, Nash SG. Psychosocial determinants of the onset and escalation of smoking: cross-sectional and prospective findings in multiethnic middle school samples. *J Adolesc Health* 2000; 27(4): 255-65.
27. Patton GC, Carlin JB, Coffey C, Wolfe R, Hibbert M, Bowes G. The course of early smoking: a population-based cohort study over three years. *Addiction* 1998; 93(8): 1251-60.
28. Hughes Y. Nicotine related disease. In: Sadock BJ, Sadock VA editors. *Kaplan and Sadock's Comprehensive text book of Psychiatry*. 7<sup>th</sup> ed. Baltimore: Lippincott Williams and Wilkins; 2000. Vol. 2. p.1033.
29. Gidwani PP, Sobol A, DeJong W, Perrin JM, Gortmaker SL. Television viewing and initiation of smoking among youth. *Pediatrics* 2002; 110(3): 505-8.